

# Fossil evidence of avian crops from the Early Cretaceous of China

Xiaoting Zheng<sup>a,b</sup>, Larry D. Martin<sup>c</sup>, Zhonghe Zhou<sup>d,1</sup>, David A. Burnham<sup>c</sup>, Fucheng Zhang<sup>d</sup>, and Desui Miao<sup>c</sup>

<sup>a</sup>Institute of Geology and Paleontology, Linyi University, Linyi, Shandong 276000, China; <sup>b</sup>Shandong Tianyu Museum of Nature, Pingyi, Shandong 273300, China; <sup>c</sup>Division of Vertebrate Paleontology, Biodiversity Institute, University of Kansas, Lawrence, KS 66045; and <sup>d</sup>Key Laboratory of Evolutionary Systematics of Vertebrates, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, China

Contributed by Zhonghe Zhou, August 3, 2011 (sent for review July 13, 2011)

**The crop is characteristic of seed-eating birds today, yet little is known about its early history despite remarkable discoveries of many Mesozoic seed-eating birds in the past decade. Here we report the discovery of some early fossil evidence for the presence of a crop in birds. Two Early Cretaceous birds, the basal ornithurine *Hongshanornis* and a basal avian *Sapeornis*, demonstrate that an essentially modern avian digestive system formed early in avian evolution. The discovery of a crop in two phylogenetically remote lineages of Early Cretaceous birds and its absence in most intervening forms indicates that it was independently acquired as a specialized seed-eating adaptation. Finally, the reduction or loss of teeth in the forms showing seed-filled crops suggests that granivory was possibly one of the factors that resulted in the reduction of teeth in early birds.**

paleornithology | feeding

Despite the discoveries of more than 30 genera of fossil birds from the Early Cretaceous lacustrine deposits in northeastern China and many examples of dietary adaptations (1), it remains unknown whether a crop was present in these birds. Recently we examined hundreds of Early Cretaceous birds housed at the Tianyu Museum of Nature in Shandong Province, China, and found several specimens belonging to two taxa with unequivocal evidence of crops. One of them is *Sapeornis*, one of the basal birds presumed to have an herbivorous diet (2, 3), and the second taxon is the basal ornithurine *Hongshanornis* (4). Both have preserved seeds in the anatomical location of the crop in extant birds. In some cases, even the soft tissue outline of the crop can be observed and resembles closely the structure in modern birds.

In extant birds a crop is a ventral pouch of the esophagus and is situated anterior to the shoulder girdle just in front of the furcula (5) (Fig. 1). In fossils its position is indicated by a roughly spherical mass of seeds that is easily discriminated from stomach contents by its location outside of the ribcage. The discovery of a crop in two phylogenetically separate avian lineages opens a window for our understanding of dietary adaptation in the early avian radiation and may also help us better understand the great diversity in the Early Cretaceous, which occurred approximately 20 million years after the earliest bird, *Archaeopteryx* (6, 7).

In modern seed-eating birds the crop provides storage, so that a number of seeds can be gathered quickly and then processed later in a more secure location without interference from competitors and/or predators. The mucus in the crop softens hard seeds so that they are more easily ground by the gizzard. The gizzard may be a more basal feature for birds, because it is widely distributed in sister groups, such as modern crocodylians. In recent birds the gizzard is posterior to the proventriculus or glandular part of the stomach. The practice of collecting large numbers of small stones in the gizzard is characteristic of seed eaters among modern birds and is often correlated with a well-developed crop.

A number of other Mesozoic birds and their theropod relatives have recently been reported with gizzard stones (gastroliths), including the basal ornithurines *Yanornis* (8) and *Archaeorhynchus* (9), the oviraptorosaur *Caudipteryx*, and the ornithomimosaur *Shenzhousaurus* (10), confirming that the presence of a muscular grinding stomach is primitive for birds. It may well have occurred in most species but is only documented in forms that foraged on the ground and thereby came into contact with small stones. The discovery of crops in combination with stomach stones demonstrates the presence at an early date of an essentially modern avian digestive system.

## Description and Comparison

Among the avian specimens we examined at the Tianyu Museum of Nature, at least three preserve clear evidence of a crop. Of these, *Sapeornis chaoyangensis* (2, 3) is the most basal bird that preserves this structure, and we discovered two examples (STM 15-15 and STM 15-29). STM 15-29 (Fig. 2 *A* and *B*) is a semi-articulated skeleton with the neck and head pulled away from the body for  $\approx 5$  cm. The skeleton is exposed dorsolaterally, with the wings folded. The right manus has moved away from the body. The sacrum is viewed dorsally, but the legs are laterally exposed. The crop is represented by a roughly circular mass ( $\approx 3$  cm in diameter) of seeds. The outside was rimmed by a layer of larger seeds ranging from 4 to 6 mm in diameter, and the anterior portion of this layer had floated out, spreading over an area of some  $9 \times 2.5$  cm (Fig. 2*B*). The center of the mass is densely packed with more than 70 smaller seeds showing an individual diameter of  $\approx 2$  mm. The seed mass was inset against the front of the furcula as in modern birds. The second specimen, STM 15-15 (Fig. 2 *C* and *D*) shows a cylindrical discoloration (darkening) of the sediments just below the ventral surface of the cervical vertebrae, extending a few centimeters anteriorly. This seems to be a trace of the esophagus, and it is faintly visible around the seed mass. The seeds were closely packed one against the other, indicating that there was little or no fleshy surface around them. The postcranial skeleton is nearly intact, with the ribcage still in its original position. Counting five vertebrae posterior to the end of the neck and within the ribcage there is a small mass of polished stones. A few of the stones are scattered away from the main concentration. The main mass indicates the position of the gizzard. Just posteroventral to this mass is a pocket of coprolitic material in the position of the intestines. The outline of the skeleton and positioning of the crop and gizzard is remarkably similar to that found in modern birds (Fig. 2).

The skeleton of *Sapeornis*, as in *Archaeopteryx*, lacks an ossified sternum, but the maxilla was edentulous, whereas the premaxilla has a small number of teeth that are elongated anteroposteriorly. The mandible was toothless and covered by a horny bill, as was

Author contributions: X.Z., L.D.M., and Z.Z. designed research; X.Z., L.D.M., Z.Z., and D.M. performed research; X.Z., D.A.B., and F.Z. contributed new reagents/analytic tools; D.A.B. and F.Z. analyzed data; and L.D.M., Z.Z., and D.M. wrote the paper.

The authors declare no conflict of interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: zhouzhonghe@ivpp.ac.cn.





